

Immediate Denture

Follow up Information

- Biting pressure on the denture will promote clotting and will decrease the initial flow of blood. However, slight bleeding can last up to 2 to 3 days.
- Use an ice compress on the affected side for 20 minutes on, and 20 minutes off for the first 36 hours.
- Limit your diet to soft nurturing foods and plenty of fluids for the next week.
- Don't take the denture out today, but rinse your mouth with warm salt water before going to bed/ Use one teaspoon of salt per cup of warm water.
- Starting tomorrow, carefully remove the denture twice a day and clean with a toothbrush and a low abrasive toothpaste or denture cleanser. Rinse your mouth with warm salt water.
- Sleep with the denture in your mouth for one week unless otherwise instructed, Then you should take it out when sleeping in order to relax the gum tissues.
- Due to the gum shrinkages that occur within the first 6 months, you may go through periods of a loose fitting denture. A temporary liner will be inserted as needed. Denture adhesives can be used during this period as well.
- Following the gum shrinkage period (about 6 months), a more permanent reline will be placed.
- You may experience sore spots caused by the denture being too long or from uneven pressure being applied to the healing gum tissues. We will adjust the denture as these problems occur.



Jeffrey A Stewart D.D.S.

FAMILY DENTISTRY • IV SEDATION
IMPLANTS • COSMETIC DENTISTRY